

1 Title: Eliciting and integrating expert knowledge to assess the viability of the critically
2 endangered golden sun-moth *Synemon plana*

3 LUIS MATA,^{1*} GEORGIA E. GARRARD,¹ ALEX S. KUTT,² BONNIE C. WINTLE,³
4 YUNG EN CHEE,^{3,4} ANNA BACKSTROM,¹ BRIAN BAINBRIDGE,⁵ JAKE URLUS,⁶
5 GEOFF W. BROWN,⁷ ARN D. TOLSMA,⁷ ALAN L. YEN,^{8,9} TIMOTHY R. NEW,¹⁰
6 SARAH A. BEKESSY,¹

7 ¹ Interdisciplinary Conservation Science Research Group, School of Global, Urban
8 and Social Studies, RMIT University, Melbourne, Vic. 3001, Australia

9 ² Australian Research Centre for Urban Ecology, Royal Botanic Gardens Victoria,
10 School of BioSciences, University of Melbourne, Parkville, Vic. 3010, Australia

11 ³ School of BioSciences, University of Melbourne, Parkville, Vic. 3010, Australia

12 ⁴ School of Ecosystem and Forest Sciences, University of Melbourne, Burnley, Vic.
13 3121, Australia

14 ⁵ Merri Creek Management Committee, 2 Lee Street, Brunswick East, Vic. 3057,
15 Australia

16 ⁶ Ecology Australia, 88B Station Street, Fairfield, Vic. 3078, Australia

17 ⁷ Arthur Rylah Institute for Environmental Research, Department of Environment,
18 Land, Water & Planning, Heidelberg, Vic. 3084, Australia

19 ⁸ Biosciences Research Branch, Department of Economic Development, Jobs,
20 Transport & Resources, La Trobe University, Bundoora, Vic. 3083, Australia

21 ⁹ School of Applied Systems Biology, La Trobe University, Bundoora, Vic. 3083,
22 Australia

23 ¹⁰ Department of Ecology, Environment and Evolution, La Trobe University,
24 Melbourne, Vic. 3086, Australia

25 *Corresponding author. RMIT University, GPO Box 2476, Melbourne, Vic. 3001,
26 Australia. Phone: +613 9925 9945. E-mail: luis.mata@rmit.edu.au

27 Running title: Golden sun-moth population viability

28 Abstract

29 The critically endangered golden sun-moth *Synemon plana* occurs in urban fringe
30 areas of southeastern Australia that are currently experiencing rapid and extensive
31 development. The urban fringe is a complex and uncertain environment in which to
32 manage threatened species with the intersection of fragmented natural habitats, built
33 environments and human populations generating novel, poorly-understood
34 interactions. In this context, management frameworks must incorporate ecological
35 processes as well as social considerations. Here we explore how biodiversity
36 sensitive urban design might improve the fate of the golden sun-moth, and
37 threatened species generally, in urban fringe environments. We: (1) developed an
38 expert-informed Bayesian Belief Network model that synthesises the current
39 understanding of key determinants of golden sun-moth population viability at sites
40 experiencing urbanising pressure; (2) quantified the nature and strength of cause-
41 effect relationships between these factors using expert knowledge; and (3) used the
42 model to assess expectations of moth population viability in response to different
43 combinations of management actions.

44 We predict that adult survival, bare ground cover and cover of resource plants are
45 the most important variables affecting the viability of golden sun-moth populations.

46 We also demonstrate the potential for biodiversity sensitive urban design as a

47 complementary measure to conventional management for this species. Our findings
48 highlight how expert knowledge may be a valuable component of conservation
49 management, especially in addressing uncertainty around conservation decisions
50 when empirical data are lacking, and how structured expert judgements become
51 critical in supporting decisions that may help ameliorate extinction risks faced by
52 threatened species in urban environments.

53 Key words: Bayesian Belief Networks, Biodiversity sensitive urban design, Grassland
54 management, Insect conservation, Threatened species management, Urban ecology

55

56 INTRODUCTION

57 The golden sun-moth *Synemon plana* Walker, 1854 (Lepidoptera, Castniidae) is a
58 listed 'critically endangered' endemic species occurring in the native grassland
59 ecosystems of southeastern mainland Australia (western and northern Melbourne,
60 and parts of the Australian Capital Territory). It is a flagship species for grassland
61 conservation, and is threatened by the severe and on-going reduction in extent of
62 native grassland habitat and the conversion of remaining grassland into degraded
63 and exotic plant dominated ecosystems (Kutt *et al.* 2015). A large proportion of the
64 moth's known distribution overlaps with urban growth areas and many populations of
65 high conservation significance now occur within a matrix of housing and industrial
66 development (Gilmore *et al.* 2008).

67 The occurrence of golden sun-moth populations adjacent to urban housing presents
68 particular challenges for habitat management, including conflicts between different
69 management actions that may be scientifically grounded but socially impractical or
70 unacceptable (Whitehead *et al.* 2014). For example, the golden sun-moth prefers
71 grasslands of low biomass that are dominated by native *Austrostipa* and
72 *Rytidosperma* that were historically maintained by native herbivore grazing and
73 periodic intense fire (Dorrough *et al.* 2004). In degraded sites, managed grazing by
74 domestic stock and controlled burning can potentially assist the species persistence
75 through the control of exotic pasture species and maintenance of low biomass
76 (O'Dwyer & Attiwill 2000). However, management by stock and fire in locations
77 adjacent to human populations is contentious because of real and perceived risks to
78 human health, lives and property (Gibbons *et al.* 2012). There is some evidence that
79 the physical structure and design of dwellings may provide habitat for known non-
80 native predators of the golden sun-moth (e.g. the common mynah *Acridotheres*

81 *tristis*) and increased predation may adversely affect golden sun-moth population
82 persistence in urban environments (Australian Government 2009).

83 Conventional management actions include measures to improve golden sun-moth
84 habitat quality through reestablishment of native grasses, weed and biomass
85 removal, and measures to reduce the mortality of golden sun-moth adults through
86 predation control. However, these 'conventional actions' may not be sufficient on
87 their own in a landscape where remnant habitats co-occur with the urban matrix.
88 Biodiversity sensitive urban design proposes a series of key principles aimed at
89 enhancing biodiversity at the site level, by improving the viability of native species
90 and ecosystems (Garrard *et al.* in review). These may involve design measures to
91 improve remnant native habitat through sympathetic management of private
92 gardens, installations that mitigate adverse impacts such as buffer zones,
93 management techniques that reduce human disturbance at important times such as
94 sanctuary periods, and initiatives to enhance human-nature interactions with
95 community engagement and education.

96 Active management is therefore an important component of sustaining golden sun-
97 moth population viability. However, there are few empirical data on cause-effect
98 linkages between the species demographic variables and conventional and
99 biodiversity sensitive urban design management actions. We therefore turn to expert
100 knowledge and knowledge engineering (Korb & Nicholson 2011) to (1) synthesise in
101 a formal model, current understanding of key determinants of golden sun-moth
102 population viability at sites experiencing urbanising pressure; (2) quantify the nature
103 and strength of cause-effect relationships between these factors using expert
104 knowledge; and (3) use the resultant model to assess expectations of golden sun-

105 moth site-level population viability, in response to different combinations of
106 management actions.

107 Our approach was driven by the need to deliver conservation-orientated
108 management solutions for a listed 'critically endangered' data-deficient Australian
109 insect species that coincides with human populations. Management actions aimed at
110 preserving the golden sun-moth in southeastern Australian peri-urban grassland
111 ecosystems may synergistically contribute to the protection of other threatened
112 grassland species (e.g. striped legless lizard *Delma impar*, matted flax-lily *Dianella*
113 *amoena*, spiny rice-flower *Pimelea spinescens spinescens*) as well as the Natural
114 Temperate Grasslands of the Victorian Volcanic Plain, which are themselves
115 critically endangered (Australian Government 2011). We are also motivated to
116 improve threatened species evaluation and policy processes by incorporating
117 structured expert opinion and exploring uncertainty, as urban landscapes are
118 undervalued and highly significant locations for such species (Ives *et al.* 2016).

119 METHODS

120 *Modelling framework*

121 We used a Bayesian Belief Network modelling framework to represent the viability of
122 golden sun-moth under different management scenarios. Bayesian Belief Networks
123 (Pearl 1988; Korb & Nicholson 2011) are graphical probabilistic models for reasoning
124 under uncertainty. Bayesian Belief Networks consist of a set of nodes that represent
125 the salient variables in the system of interest. Uncertainty is represented by
126 specifying probability distributions for the node variables (which can be continuous or
127 discrete). Arcs (or arrows) indicate where conditional dependencies exist between
128 'parent' (denoted $pa(X)$) and 'child' (denoted $P(X)$) nodes. For each variable, all

129 relevant (and mutually exclusive states) are defined. Each child node has a
130 conditional probability table that quantifies the probabilistic effects that parent nodes
131 have on it, that is, $P(X|pa(X))$.

132 The graphical network of nodes and arcs expresses the chain of logic or causal
133 argument that links variables to outcomes. When the graphical structure is fully
134 specified, the conditional probability tables parameterised, and the Bayesian Belief
135 Network is compiled, it can be used for predictive reasoning about the system. Users
136 can set the values of any combination of nodes in the network. This 'evidence', e ,
137 propagates through the network producing a new posterior probability distribution
138 ($P(X|e)$) for each node in the network. In the Bayesian Belief Network modelling
139 software that we use (Netica, version 5.18, Norsys Software Corporation), a number
140 of efficient exact and approximate inference algorithms are available for performing
141 this belief updating. A particular benefit of Bayesian Belief Networks is that
142 knowledge and data from multiple sources such as theoretical insight, empirical data,
143 output from statistical or process models and expert judgements can be used to
144 construct the graphical structure and parameterise the conditional probability tables
145 (Cain 2001).

146 *Model development*

147 The goal was to capture the key factors that influence the population viability of
148 golden sun-moth at sites experiencing urbanising pressure. The three main tasks are
149 selection and definition of variables, specification of the Bayesian Belief Network
150 graphical structure (i.e. network of nodes and arcs) and construction of conditional
151 probability tables for each node. We developed a first-cut Bayesian Belief Network

152 using a review of the literature. We then used an expert workshop to revise the
153 model and parameterise the conditional probability tables for each node.

154 We searched for 'golden sun-moth' and 'golden sun moth' in the Web of Knowledge,
155 Scopus and Google Scholar (April 2014). From this literature, we identified key
156 variables that influence the population viability of golden sun-moth and their putative
157 cause-and-effect relationships. We also incorporated five biodiversity sensitive urban
158 design features, namely 'Ecological buffer zone', 'Fire buffer zone', 'Clean
159 construction', 'Viewing platforms' and 'Sanctuary periods', as we wanted to
160 investigate their influence on golden sun-moth population viability. These features
161 were chosen as they adhere to the key principles for biodiversity sensitive urban
162 design (i.e. maintain or introduce habitat, facilitate dispersal, minimise threats and
163 anthropogenic disturbances, facilitate natural ecological processes, and facilitate
164 positive human-nature interactions). Some of these have also been previously
165 assessed by Garrard *et al.* (in review) in a study involving the persistence of the
166 native temperate grasslands of the Victorian Volcanic Plain. The first-cut Bayesian
167 Belief Network and the literature used to develop it are given in Figure A1
168 (Supporting Information). In developing this literature-based Bayesian Belief
169 Network, we took care to apply the following recommendations given in Marcot *et al.*
170 (2006), Korb & Nicholson (2011) and Chen & Pollino (2012): (1) the number of
171 parent nodes to any given child node was kept to three or less; (2) a balance
172 between parsimony and precision was sought when deciding on the number of
173 necessary discrete states within each node; and (3) continuous correlates were
174 discretised as appropriate. As the joint probabilistic effects of parents on child nodes
175 were to be assessed by experts, these guidelines help to ensure that the structure
176 did not impose a heavy cognitive burden on the assessment task.

177 We refined the literature-based Bayesian Belief Network and populated the
178 conditional probability tables in a one-day workshop involving five specialists with
179 expertise in golden sun-moth ecology and conservation, in July 2014. The experts
180 included an academic with decades of entomological research experience, a
181 research entomologist at a leading government agency and environmental
182 consultants with extensive field experience in golden sun-moth survey protocols. All
183 experts had authored one or more peer-reviewed publications and/or reports in
184 which the main focus of research had been the golden sun-moth. Prior to the
185 workshop, the experts received a training document, in which they were provided
186 with information on the facilitators, the workshop's goals, biodiversity sensitive urban
187 design principles and expert elicitation methodologies.

188 During the workshop, the five experts (ASK, ALY, BB, JU & TRN), supported by
189 three facilitators (BCW, GEG & LM), established the spatial and temporal context for
190 the Bayesian Belief Network model that would be built. It was agreed that the model
191 would focus on grassland patches of 10-20 hectares, located in areas about to be
192 disturbed by urban development. The model timeframe was set to 1-3 years, since it
193 is presently unclear whether the golden sun-moth life cycle takes one, two or three
194 years to complete (New 2012). An agreement was also reached to work exclusively
195 with input variables that could be modified through management. Consequently,
196 environmental variables such as temperature, though important, were excluded.
197 Instead, we assumed 'average' temperature conditions for the modelling exercise.

198 After agreeing on the modelling context, the experts were given a detailed model
199 walkthrough of the literature-based Bayesian Belief Network. This formed the starting
200 point for discussions about candidate output, intermediate and input variables;
201 exactly what each represented, how they ought to be described and defined, and

202 how they related to any parent or child variables. Using the Bayesian Belief Network
203 modelling software, Netica (version 5.18, Norsys Software Corporation),
204 modifications to the model's structure were incorporated and removed dynamically
205 by the facilitators as the discussion proceeded. After multiple rounds of discussions,
206 experts and facilitators agreed on a consensus Bayesian Belief Network that they felt
207 was a good representation of current knowledge about key influences on the
208 population viability of the golden sun-moth.

209 *Parameterisation of the peer-reviewed Bayesian Belief Network using expert*
210 *knowledge*

211 The strength of the relationships between conditionally dependent variables in the
212 graphical model was assessed and parameterised using expert elicitation. We
213 followed the guidelines provided in Kuhnert *et al.* (2010), Martin *et al.* (2012) and
214 McBride & Burgman (2012) to design the process by which knowledge was elicited
215 from the experts. Prior to running the elicitation to parameterise the golden sun-moth
216 model, the experts completed a practice run to familiarise themselves with the task
217 of conditional probability table assessment. We also used a percentage scale (0-
218 100) rather than a probability scale (0-1), as research suggests that people find
219 probabilities difficult to understand and reason with (Gigerenzer & Hoffrage 1995).
220 Each expert completed all the conditional probability tables in the model
221 independently and privately, resulting in five parameterised Bayesian Belief
222 Networks. We also created a combined consensus model by pooling individual node
223 conditional probability table judgements through simple averaging.

224 *Model evaluation*

225 The individual expert models as well as the final combined model were evaluated
226 using two types of sensitivity analysis: sensitivity to evidence and sensitivity to
227 changes in parameters. Sensitivity to evidence tells us how much a finding at one
228 node will likely change the beliefs at another (the so-called 'query' node). We used
229 this to identify which variables have the greatest influence on the output node
230 'Change in golden sun-moth population'. In Netica, the 'sensitivity to findings'
231 function uses entropy reduction (measured in bits) to measure the effect of one
232 variable on another. The greater the entropy reduction value associated with a
233 findings node, the greater its influence on the query node.

234 In this study, the outcome of greatest concern was when the 'Change in golden sun-
235 moth population' variable, was in the state 'Decline'. We therefore conducted our
236 sensitivity to changes in parameters analysis with specific reference to this outcome.
237 This involved noting the posterior probability of this outcome, as the state of each
238 node in the Bayesian Belief Network was altered between its minimum and
239 maximum range (Pollino *et al.* 2007, Korb & Nicholson 2011). This analysis can tell
240 us for which variables, greater precision in estimation would be useful.

241 Finally we also undertook scenario-based evaluation to examine the expected
242 'Change in golden sun-moth population' associated with a series of scenarios of
243 management interest.

244 The .neta extension 'Netica Bayesian Belief Network' files containing the necessary
245 expert-parameterised conditional probability tables to re-run the analyses are
246 provided in the online Supporting Information.

247 *External review*

248 As a means of further evaluating the consensus model, we sought external peer-
249 review (Marcot *et al.* 2006). We asked the experts who had participated in the
250 workshop to recommend other suitably qualified golden sun-moth experts. Of the
251 recommended experts who were contacted, three agreed to assist with the external
252 peer-review. Either in person or via videoconference, we stepped each expert
253 individually through the process that led to the consensus model. We asked the
254 experts for specific feedback on whether: (1) the model variable names and states
255 were appropriately and adequately defined with respect to the spatial and temporal
256 scale and specific problem context; (2) the overall graphical structure of the model
257 was based on sound ecological reasoning; and (3) all important variables had been
258 included in the model and whether any omissions were justifiable/defensible. The
259 external reviewers were further requested to provide a 'reasonableness' check on
260 node relationships encoded in the conditional probability tables. The external
261 reviewers were provided with all workshop outputs, including the 'Netica Bayesian
262 Belief Network' (.neta) files necessary to re-run the analyses. Of the three experts
263 who were briefed to conduct the external review, two provided feedback (ADT &
264 GWB).

265 RESULTS

266 The consensus Bayesian Belief Network model is composed of 14 nodes and 16
267 arcs (Fig. 1), and the names, states, descriptions and explanations of all model
268 variables are summarised in Table 1. The graphical model is structured according to
269 the main conceptual ideas as follows:

- 270 1. The viability of golden sun-moth at urban fringe sites is believed to be strongly
271 linked to the magnitude of 'Change in golden sun-moth population' over a 1-3
272 year timeframe.
- 273 2. The golden sun-moth population includes short-lived adults and larval stages
274 of variable longevity. In the model therefore, 'Change in golden sun-moth
275 population' depends explicitly on 'Adult survival', while the contribution of
276 larval golden sun-moth stages is represented indirectly by 'Cover of resource
277 plants' and 'Bare ground cover' which both influence the survival of the larval
278 stages.
- 279 3. 'Adult survival' is affected by whether 'Predation management' is implemented
280 or not. 'Cover of resource plants' depends on whether native grasses are re-
281 established and how much weed cover there is in the grassland patch. 'Bare
282 ground cover' which is important for the larval stages is determined by 'Weed
283 cover' and 'Biomass management type'.
- 284 4. 'Weed cover' in turn, is driven by the strength of the 'Weed propagule
285 pressure', the amount of nitrogen and phosphorus reaching the grassland
286 patch ('Soil inputs'), and whether 'Weed management' follows standard
287 practice or is absent.
- 288 5. Construction practices during development ('Clean construction') have an
289 impact on 'Weed propagule pressure', as does the type of 'Buffer zone'. In
290 addition, the type of 'Buffer zone' influences nitrogen and phosphorous inputs
291 to the site and constrains the 'Biomass management type' that can be applied
292 (e.g. burning to remove excess biomass is infeasible in the absence of a
293 buffer zone between built environments and a grassland patch).

294 6. 'Community engagement' based around informed discussion of benefits and
295 risks of biomass management options is expected to increase the
296 acceptability of burning as a tool.

297 Entropy reduction values calculated in the sensitivity to evidence analysis allowed us
298 to produce a ranking of the network variables, in order of influence on the 'Change in
299 golden sun-moth population' query node (Table 2, Table A1 in the online Supporting
300 Information).

301 Though there were slight differences in the variables ranked from 2 to 13, experts
302 were largely and consistently in agreement about the relative importance of
303 variables. In the combined model, as well as for each expert-parameterised model,
304 'Adult survival' was the variable that most influenced golden sun-moth viability. The
305 sensitivity analysis indicated that changes to the golden sun-moth population were
306 most influenced by its parent nodes 'Adult survival', 'Cover of resource plants' and
307 'Bare ground cover', and least sensitive to the most distal nodes such as the
308 biodiversity sensitive urban design input nodes 'Community engagement', 'Clean
309 construction', and type of 'Buffer zone' (Fig. 2). This is not surprising and these
310 results reflect the logic represented by the graphical structure of the network.

311 Using the combined Bayesian Belief Network, we examined multiple scenarios to
312 probe the expected response of golden sun-moth to different sets of management
313 actions. As a basic check, we corroborated that setting the three most influential
314 network variables of 'Adult survival', 'Bare ground cover' and 'Cover of resource
315 plants' to their lowest value shifted the probability mass of 'Change in golden sun-
316 moth population' strongly to the 'Decline' state. By contrast, a shift in the opposite

317 direction occurred when these variables were set to their highest values (Table 3 and
318 Fig. 3).

319 When the full suite of conventional management options of 'Predation management',
320 'Weed management' and 'Reestablishment of native grasses' were all set to their
321 highest values, the most likely state of 'Change in golden sun-moth population' was
322 'Stable' (Conventional Management *best-case scenario* in Table 3). In contrast,
323 when these options were at their lowest values, the most likely state of 'Change in
324 golden sun-moth population' was 'Decline' (Conventional management *worst-case*
325 *scenario* in Table 3).

326 There is a small difference in the expected outcome when the biodiversity sensitive
327 urban design variables (i.e. 'Clean construction', 'Buffer zone' and 'Community
328 engagement') were set to their highest or lowest values (Biodiversity sensitive urban
329 design *best-case scenario* and biodiversity sensitive urban design *worst-case*
330 *scenario* in Table 3). In both scenarios, the most likely state is 'Stable', and the
331 difference in probabilities for each of the states 'Decline' and 'Increase' (between the
332 two scenarios) was approximately 3%. Enacting all biodiversity sensitive urban
333 design options in addition to conventional options (i.e. Conventional management +
334 Biodiversity sensitive urban design *best-case scenario*) demonstrated some support
335 for this management approach, with the probability of the 'Increase' state of 'Change
336 in golden sun-moth population' increasing from 31.7 to 33.0% (Table 3).

337 DISCUSSION

338 The results of our study suggest that adult survival, bare ground cover and cover of
339 resource plants are the most important variables affecting the viability of golden sun-
340 moth populations, and this corresponds to field evidence for the species collected

341 across its range (O'Dwyer and Attiwill 1999; Brown *et al.* 2012; Richter *et al.* 2013a).
342 In addition, outputs from the scenario-based evaluations further suggest that a best-
343 case scenario in which all three variables are simultaneously tested at their higher
344 states has the potential to improve golden sun-moth populations from a stable to
345 increasing state (i.e. change in population size greater than 25%). By contrast, a
346 worst-case scenario in which these variables are tested at their lowest states is
347 predicted to change the state of golden sun-moth populations from stable to
348 declining (i.e. change in population size greater than -25%). Taken together, these
349 findings highlight the interacting and pivotal role that management of adult
350 survivorship, ground cover and food resources have for the conservation of this
351 species. Actions that are designed to optimise the state of these key population and
352 habitat variables are predicted to enhance the persistence of golden sun-moth
353 populations into the future.

354 When the model variables were assessed individually, our results show adult
355 survival to be the most influential variable affecting golden sun-moth population
356 viability. Our model further identified predation management as the single-most
357 important controllable variable influencing adult survival. Management of introduced
358 predators is particular important given that naturally co-occurring species, such as
359 the striped legless lizard *Delma impar* (Kutt *et al.* 1998), also prey on golden sun-
360 moths. These findings suggest that management and urban design that a) minimises
361 the degradation of native vegetation and b) reduces human-made structures that
362 could facilitate species predation on the golden sun-moth, are key. The Australian
363 Government impact assessment guidelines for the golden sun-moth, for example,
364 indicate that moth predation by insectivorous birds (e.g. willie wagtail *Rhipidura*
365 *leucophrys*) may be avoided or mitigated by limiting the availability of nesting and

366 breeding structures and by designing fences that allow passage of adult golden sun-
367 moth while simultaneously limiting perching surfaces (Australian Government 2009).

368 The cover of resource plants was the second most influential variable affecting the
369 population viability of the golden sun-moth. This variable in turn was most strongly
370 affected by the extent of weed cover and the implementation of management actions
371 aimed at the re-establishment of native grasses. Understanding of the full range of
372 consumable plants for larval golden sun-moth, and of the optimal density, condition
373 and species of these, is still very incomplete – but the critical importance of larval
374 food plants in site restoration to support and enhance golden sun-moth populations
375 underpins practical conservation management for the moth. Threshold density of a
376 key host plant, *Rytidosperma erianthum*, was assessed experimentally at Mt. Piper
377 (Broadford, Victoria) by combining weeding with the planting of seedlings (O'Dwyer &
378 Attiwill 2000). The elimination of competition from weeds provided significant benefit,
379 and sites with golden sun-moth had *Rytidosperma* cover of >40 %, a level
380 subsequently cited as a target threshold for site quality.

381 A major alien invader of grassland sites, Chilean needle grass *Nassella neesiana*, is
382 a declared noxious weed – with an obligation to eradicate it wherever it is found. It
383 occurs on many grasslands occupied by golden sun-moth, and large moth
384 populations have been found on grassland patches comprised entirely of *Nassella*
385 (Richter *et al.* 2013a). Pupal case surveys implied a close association with the weed,
386 endorsing earlier suppositions (Gilmore *et al.* 2008; Brown *et al.* 2012) that *Nassella*
387 may be a supplementary or primary food for golden sun-moth larvae in Victoria,
388 where the native grass species have been reduced or eliminated. This presents a
389 clear conservation dilemma, the conflict between the legal requirement to eliminate
390 or prevent the spread of a declared noxious weed and its potential role as a key food

391 source for a critically endangered moth species in degraded grassland patches in
392 which alternative, native, food plants are sparse. The relative priority of weed control
393 versus golden sun-moth population management should be context-specific for each
394 individual grassland patch.

395 Bare ground cover was found to be the third most important variable influencing
396 golden sun-moth population viability, and this variable was directly affected by weed
397 cover and biomass management type. The golden sun-moth prefers an open
398 tussock structure with sparse inter-tussock spaces (O'Dwyer & Attiwill 2000; Gilmore
399 *et al.* 2008; Australian Government 2009), and patches of bare ground may be
400 important during various stages of their lifecycle, especially reproduction. Females
401 are semi-flightless and, after emerging from the pupa, they tend to remain on the
402 ground, flashing their brightly-coloured hindwings from a conspicuous location to
403 attract low-flying patrolling males (Australian Government 2009). Areas of bare
404 ground, often covered by bryophytes, may also be an indication of native grasslands
405 in good condition (Australian Government 2011). For example, *Themeda*-dominated
406 grasslands without appropriate biomass control may form a thick thatch of vegetation
407 that chokes out other native species (Morgan & Lunt 1999). With biomass reduction,
408 competitive exclusion may be prevented, allowing the growth of grasses preferred by
409 golden sun-moth, such as *Austrostipa* and *Rytidosperma*. Grasslands of low
410 biomass and dominated by golden sun-moth preferred grasses were historically
411 maintained by grazing by native herbivores and periodic fire (Dorrrough *et al.* 2004),
412 and such 'natural' disturbance would be considered optimal for controlling biomass.
413 In degraded sites, controlled grazing by domestic stock has assisted in the control of
414 exotic pasture species (O'Dwyer & Attiwill 2000). However, grazing by heavy stock
415 can lead to increased soil compaction and decreased water infiltration, and soils in

416 pastures that are even lightly grazed may eventually reach the same compacted
417 state as heavily-grazed pastures (Greenwood & McKenzie 2001).

418 The contentious and difficult social problems created by grazing and fire
419 management actions in locations adjacent to human populations, including the
420 potential for loss of property (Gibbons *et al.* 2012), has led to suggestions that other
421 management solutions such as slashing, mowing and weed spraying to control grass
422 biomass and weed species might be more appropriate in an urban setting (Australian
423 Government 2009). However, land managers need to recognise the potential
424 impacts of these alternative solutions. For example, compressive and
425 sliding/shearing forces by the wheels of agricultural vehicles, particularly when soils
426 are damp, are principal causes of soil compaction (Batey 2009). Much also remains
427 to be learnt about the effects of herbicides in natural ecosystems, particularly their
428 impacts on insects and other invertebrates (Pratt *et al.* 1997).

429 Our results suggest that amongst the variables included in our model, those
430 representing biodiversity sensitive urban design (i.e. clean construction, an
431 appropriate buffer zone and community engagement) were individually and
432 collectively unlikely to exert a large influence over the viability of golden sun-moth
433 populations. This was anticipated, as these variables are indirect actions, located
434 furthest from the output node in the model. However, we recommend a cautious
435 approach to interpreting these findings. Biodiversity sensitive urban design aims to
436 mitigate the severe impacts of urbanisation on biodiversity by improving the *in situ*
437 viability of native species and ecosystems (Garrard *et al.* in review). This is in
438 contrast to common approaches to compensate for biodiversity and habitat losses in
439 urban areas via off-site offsets. Offsetting is unlikely to achieve net positive
440 outcomes for biodiversity (Bekessy *et al.* 2010), particularly in the case of critically

441 endangered ecosystems where available offset sites are limited (Gordon *et al.* 2011).
442 The assessment of the influence of biodiversity sensitive urban design on species
443 viability requires the integration of social and ecological variables, for example, to
444 determine in this case the extent to which engagement with the community may
445 indirectly influence bare ground cover by improving understanding of and support for
446 specific biomass control measures such as fuel reduction burns. Existing research
447 and evidence for these relationships is scarce, even when compared to the paucity
448 of ecological information for data-deficient species like the golden sun-moth.
449 Perhaps either our model or the domain experts that parameterised it was/were
450 overly cautious about the potential benefits of biodiversity sensitive urban design.
451 The potential of biodiversity sensitive urban design actions to mitigate *in situ* the
452 detrimental impacts of poorly-planned urban development remains to be fully
453 empirically tested.

454 The results from our study highlight how expert knowledge may be a valuable
455 component of conservation management, especially in addressing uncertainty
456 around conservation decisions when empirical data are lacking. However, it is
457 important to acknowledge that our model is a literature-based, expert-judged
458 approximation of the causal web of key correlates affecting the population viability of
459 the golden sun-moth, and expert judgements are not without their biases. While a
460 group of experts is likely to be less biased than any given individual (Burgman *et al.*
461 2011), experts within a narrow domain are not wholly independent from each other,
462 because they tend to source knowledge from a similar literature, and often share
463 similar beliefs. Arguably, a more accurate representation could be achieved by
464 generating the model using data derived from empirical studies; however, in our
465 case, few empirical data exists. Therefore, when pressing conservation actions are

466 warranted and empirical data are lacking, structured expert judgements become
467 critical in supporting decisions that may help ameliorate extinction risks faced by
468 threatened species. When experts make judgements within their domain of
469 expertise, and when those judgements are elicited and aggregated in a transparent
470 and repeatable way using approaches that mitigate common biases such as group
471 think and halo effects, their judgements are almost certainly better than the
472 alternative: i.e. relying on no evidence or opaque and unstructured lay estimates to
473 make decisions (Aspinall 2010, Burgman et al. 2011). If poorly formulated, those
474 decisions could have strong detrimental impacts on the focal species.

475 There were several important limitations of the study including issues raised by the
476 external experts who reviewed the model. We examined the causal web of key
477 correlates affecting the viability of golden sun-moth populations in city fringe
478 grasslands prior to hypothetical disturbance by urban development, and hence our
479 results do not necessarily extend beyond this particular context. For example, our
480 modelling approach does not incorporate variables that are beyond our control,
481 notably abiotic environmental variables such as temperature and precipitation.
482 Focusing exclusively on drivers of golden sun-moth viability for which there are
483 potential management solutions might be an issue, most notably when the golden
484 sun-moth is expected to do poorly as a consequence of the strong influence of an
485 abiotic environmental factor (e.g. extreme temperatures). The challenge remaining,
486 then, will be to incorporate the interactive effects of abiotic environmental and
487 management variables to better understand the effect of the former on the efficacy of
488 the latter.

489 Bayesian Belief Networks can also be extended to explicitly aid decision-making by
490 including decision nodes to represent specific choices and utility nodes to measure

491 the cost of particular choices as well as the value of predicted outcomes. Future
492 investigations could benefit from addressing other important issues relevant to
493 golden sun-moth conservation such as habitat contraction and degradation outside
494 urban environments, habitat heterogeneity and its influence on population fluctuation
495 and survivorship (Kutt *et al.* 2015) and the potential impacts of climate change.
496 Although we strived to elucidate and include all relevant variables and links (given
497 the constraints to minimise probability elicitation and unnecessary uncertainty
498 propagation), the external experts considered that the model may have been more
499 informative if it had included further management impacts such as compaction and
500 spraying. Other variables, such as 'commercial or government investment' or 'land
501 acquisition', would also be good to include in future iterations of the model. The
502 external reviewers indicated that 'biomass management' (i.e. slashing, grazing and
503 burning) may warrant separation into distinct actions, since the effect of each
504 approach on structure and floristics may vary, and all three methods would not be
505 used together in one location, or at least not simultaneously.

506 CONCLUSIONS

507 The golden sun-moth is a critically endangered species that occurs in urban fringe
508 areas that will experience substantial future development. Management of this
509 species requires tools to help make sound conservation management and planning
510 decisions in the face of complex socio-ecological processes and substantial
511 uncertainty. Our findings are relevant at multiple levels. First, our results may be
512 applied to the management of the golden sun-moth in urban environments as they
513 indicate the important role of adult survival, bare soil cover and cover of resource
514 plants for the population viability of the species. Second, we investigated the
515 potential of biodiversity sensitive urban design as a complementary measure to

516 conventional management for this species, and demonstrate some support for this
517 approach; the *in-situ* nature of this approach contrasts with typical urban design
518 scenarios that seek to offset biodiversity from areas to be developed and forgo
519 onsite values. Finally, our study provides a good example of structured elicitation
520 and aggregation of expert knowledge to address uncertainty around conservation
521 decisions when ecological data are lacking.

522 ACKNOWLEDGEMENTS

523 This research is supported by the Australian Government's National Environmental
524 Science Programme Clean Air and Urban Landscapes Hub (NESP CAUL), the
525 Australian Research Council Centre of Excellence for Environmental Decisions (ARC
526 CEED) and The Myer Foundation. Sarah Bekessy is supported by an Australian
527 Research Council Future Fellowship. We are grateful to Libby Rumpff for generously
528 providing help with the Bayesian Belief Networks, and to two independent reviewers
529 for providing valuable feedback on an earlier version of the manuscript.

530 REFERENCES

531 Aspinall W.A. (2010) A route to more tractable expert advice. *Nature* **243**, 294–95.

532 Australian Government (2009) Significant impact guidelines for the critically
533 endangered golden sun moth (*Synemon plana*). Commonwealth of Australia.

534 Australian Government (2011) Nationally Threatened Ecological Communities of the
535 Victorian Volcanic Plain: Natural Temperate Grassland and Grassy Eucalypt
536 Woodland. Commonwealth of Australia.

537 Batey T. (2009) Soil compaction and soil management – a review. *Soil Use Manage.*
538 **25**, 335–45.

539 Bekessy S.A., Wintle B.A., Lindenmayer D.B., McCarthy M.A., Colyvan M., Burgman
540 M.A., Possingham H.P. (2010) The biodiversity bank cannot be a lending bank.
541 *Conserv. Lett.* **3**, 151–8.

542 Brown G., Tolsma A., McNabb E. (2012) Ecological aspects of new populations of
543 the threatened Golden sun moth *Synemon plana* on the Victorian Volcanic Plains.
544 *The Victorian Nat.* **129**, 77–85.

545 Burgman M.A., McBride M.L., Ashton R., Speirs-Bridge A., Flander L., Wintle B.C.,
546 Fidler F., Rumpff L. & Twardy C. (2011) Expert status and performance. *PLoS ONE*
547 **6**, e22998.

548 Cain J. (2001) *Planning improvements in natural resources management –*
549 *Guidelines for using Bayesian networks to support the planning and management of*
550 *development programmes in the water sector and beyond*. Centre for Ecology and
551 Hydrology, Wallingford, United Kingdom.

552 Chen S.H. & Pollino C.A. (2012) Good practice in Bayesian network modelling.
553 *Environ. Model. Softw.* **37**, 134–45.

554 Dorrough J., Yen A., Turner V., Clark S. G., Crosthwaite J. & Hirth J. R. (2004)
555 Livestock grazing management and biodiversity conservation in Australian temperate
556 grassy landscapes. *Aust. J. Agric. Res.* **55**, 279–95.

557 Gibbons P., van Bommel L., Gill A.M., Cary G.J., Driscoll D.A., Bradstock R.A.,
558 Knight E., Moritz M.A., Stephens S.L. & Lindenmayer D.B. (2012) Land management
559 practices associated with house loss in wildfires. *PLoS ONE* **7**, e29212.

560 Gilmore D., Koehler S., O'Dwyer C. & Moore W. (2008) Golden Sun Moth *Synemon*
561 *plana* (Lepidoptera: Castniidae): results of a broad survey of populations around
562 Melbourne. *The Victorian Nat.* **125**, 39–46.

563 Gigerenzer G. & Hoffrage U. (1995) How to improve Bayesian reasoning without
564 instruction – Frequency formats. *Psychol. Rev.* **102**, 684–704.

565 Gordon A., Langford W.T., Todd J.A., White M.D., Mullerworth D.W., Bekessy S.A.
566 (2011) Assessing the impacts of biodiversity offset policies. *Environ. Model. Softw.*
567 **26**, 1481–8.

568 Greenwood, K.L. & McKenzie B.M. (2001). Grazing effects on soil physical
569 properties and the consequences for pastures: a review. *Aust. J. Exp. Agric.* **41**,
570 1231–50.

571 Ives C.D., Lentini P.E., Threlfall C.G., Ikin K., Shanahan D.F., Garrard G.E., Bekessy
572 S.A., Fuller R.A., Mumaw L., Rayner L., Rowe R., Valentine L.E. & Kendal D. (2016)
573 Cities are hotspots for threatened species. *Glob. Ecol. Biogeogr.* **25**, 117–26.

574 Korb K.B. & Nicholson A.E. (2011) *Bayesian artificial intelligence*. CRC Press, Boca
575 Raton, USA.

576 Kuhnert P.M., Martin T.G. & Griffiths S.P. (2010) A guide to eliciting and using expert
577 knowledge in Bayesian ecological models. *Ecol. Lett.* **13**, 900–14.

578 Kutt A.S., Coulson G. & Wainer J. (1998) Diet of the striped legless lizard *Delma*
579 *impar* (Squamata: Pygopodidae) in a Western (Basalt) Plains Grassland, Victoria.
580 *Aust. Zool.* **31**, 13–9.

581 Kutt A.S., McKenzie V.J., Wills T.J., Retallick R.W.R., Dalton K., Kay N. & Melero-
582 Blanca E. (2015) Spatial and temporal determinants of golden sun moth *Synemon*
583 *plana* distribution. *Austral Ecol.* **40**, 100–7.

584 Marcot B.G., Steventon J.D., Sutherland G.D. & McCann R.K. (2006) Guidelines for
585 developing and updating Bayesian belief networks applied to ecological modeling
586 and conservation. *Can. J. For. Res.* **36**, 3063–74.

587 Martin T.G., Burgman M.A., Fidler F., Kuhnert P.M., Low-Choy S., McBride M.F. &
588 Mengersen K. (2012) Eliciting expert knowledge in conservation science. *Conserv.*
589 *Biol.* **26**, 29–38.

590 McBride M.F. & Burgman M.A. (2012) What is expert knowledge, how is such
591 knowledge gathered, and how do we use it to address questions in landscape
592 ecology? In: *Expert knowledge and its application in landscape ecology* (eds. A.H.
593 Perera *et al.*). Springer Science + Business Media.

594 Morgan J.W. & Lunt I.D. (1999). Effects of time-since-fire on the tussock dynamics of
595 a dominant grass (*Themeda triandra*) in a temperate Australian grassland. *Biol.*
596 *Conserv.* **88**, 379–86.

597 New T.R. (2012) The Golden sun moth, *Synemon plana* Walker (Castniidae):
598 continuing conservation ambiguity in Victoria. *The Victorian Nat.* **129**, 109–13.

599 New T.R., Gibson L. & Van Praagh B.D. (2007) The Golden Sun-moth *Synemon*
600 *plana* (Castniidae) on Victoria's remnant southern native grasslands. *The Victorian*
601 *Nat.* **124**, 254–7.

602 O'Dwyer C. & Attiwill P.M. (1999) A comparative study of habitats of the Golden Sun
603 Moth *Synemon plana* Walker (Lepidoptera: Castniidae): implications for restoration.
604 *Biol. Conserv.* **89**, 131–41.

605 O'Dwyer C. & Attiwill P. M. (2000) Restoration of a native grassland as habitat for the
606 Golden Sun Moth *Synemon plana* Walker (Lepidoptera; Castniidae) at Mount Piper,
607 Australia. *Restor. Ecol.* **8**, 170–4.

608 Pearl J. (1988) *Probabilistic reasoning in intelligent systems: networks of plausible*
609 *inference*. 2nd edition. Morgan Kaufmann Publishers, San Francisco.

610 Pollino C.A., Woodberry O., Ann Nicholson A., Korb K. & Hart B.T. (2007)
611 Parameterisation and evaluation of a Bayesian network for use in an ecological risk
612 assessment. *Environ. Model. Softw.* **22**, 1140–52.

613 Pratt, J.R., Melendez, A.E., Barreiro, R., and Bowers, N.J. (1997). Predicting the
614 ecological effects of herbicides. *Ecol. Appl.* **7**, 1117–24.

615 Richter A., Osborne W., Hnatiuk S. & Rowell A. (2013a) Moths in fragments: insights
616 into the biology and ecology of the Australian endangered golden sun moth
617 *Synemon plana* (Lepidoptera: Castniidae) in natural temperate and exotic grassland
618 remnants. *J. Insect Conserv.* **17**, 1093–104.

619 Richter A., Weinhold D., Robertson G., Young M., Edwards T., Hnatiuk S. & Osborne
620 W. (2013b) More than an empty case: a non invasive technique for monitoring the
621 Australian critically endangered golden sun moth, *Synemon plana* (Lepidoptera:
622 Castniidae). *J. Insect Conserv.* **17**, 529–36.

623 Whitehead A.L., Kujala H., Ives C.D., Gordon A., Lentini P.E., Wintle B.A., Nicholson
624 E. & Raymond C.M. (2014) Integrating biological and social values when prioritizing
625 places for biodiversity conservation. *Conserv. Biol.* **28**, 992–1003.

Table 1 Nodes of the final peer-reviewed Bayesian Belief Network representing the causal web of key correlates affecting the population viability of the golden sun-moth *Synemon plana* in southeastern Australian peri-urban grassland ecosystems.

Node	Type	States	Description	Importance
Output node				
Change in golden sun-moth population	C	Decline (> -25%) Stable (-25% – 25%) Increase (> 25%)	Percentage inter-annual variation in population size.	A strong indicator of the viability of the population.
Intermediate nodes				
Adult survival	D	Below average Average Above average	Probability that adult individuals will survive at least into reproductive stage.	Linked to higher rates of female oviposition.
Bare ground cover	C	Low (< 15%) Average (15 – 25%) High (> 25%)	Percentage of grassland not covered by vegetation.	The species immature stages develop in the ground.
Biomass management type	D	Absent Slashing Grazing Burning	Method used to manage the grassland's excess biomass.	Removal of excess biomass prevents sprouting of non-resource plant species.
Cover of resource plants	C	Low (< 10%) Average (10 – 30%) High (> 30%)	Percentage of grassland covered by resource plant species.	Resource plants are critical for the species to feed and reproduce.
Soil inputs	D	Low Average High	Amount of external nitrogen and phosphorous reaching the grassland.	Higher rates of nitrogen and phosphorous will favour weed establishment.
Weed cover	C	Low (< 15%) Average (15 – 75%) High (> 75%)	Percentage of grassland covered by weed plant species.	Weeds compete directly with resource plants whilst reducing bare ground cover.
Weed propagule pressure	D	Low Average High	Amount of weed seeds reaching the grassland from adjacent areas.	Linked to higher rates of weed establishment.
Input nodes				
Conventional management actions				
Predation management	D	Not implemented Implemented	Actions taken to prevent adult mortality by	Preventing predation is associated with

			non-native predators.	higher rates of ⁶²⁶ adult survival.
Reestablishment of native grasses	D	Not implemented Implemented	Actions taken to increase the amount of native grasses present in the grassland.	Reestablishing native grasses ⁶²⁷ is associated with higher densities of resource plants.
Weed management	D	Absent Standard	Actions taken to remove weeds from the grassland.	Weed removal is linked to a decrease in weed cover.
<i>Biodiversity sensitive urban design actions</i>				
Buffer zone	D	Absent Impervious Pervious	Establishment of either impervious (e.g., street) or pervious (e.g., vegetated) surface buffering the grassland from the development.	A buffer zone may reduce the likelihood of nutrient runoff spilling from the developed area into the grassland.
Clean construction	D	Not implemented Implemented	Actions taken to minimise the introductions of weed seeds during development.	Linked to a decrease in weed propagule pressure.
Community engagement	D	Absent Present	Actions taken to educate the community on the pros and cons of conventional biomass management actions.	Associated with an increase likelihood of accepting burning as a safe option to managed the grassland's excess biomass.
C: Continuous; D: Discrete				